

CTNNB1 Knockdown 293T Cell Lysate, Heterozygous

Catalog No.: RM50085

Basic Information

Catalog No.

RM50085

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockdown

Gene Information

Gene Symbol

CTNNB1

Species

Human

Gene ID

1499

Swiss Prot

P35222

SynonymsEVR7; CTNNB; MRD19; NEDSDV;
armadillo; in

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.

Product Information

Description

CTNNB1 Knockdown cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:1bp insertion in exon2

Allele-2:WT

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CATGGAACCAAGACAGAAAA GCGGCTGTTAGTCACTGGC
Mut CATGGAACCAAGACAGAAAAAGCGGCTGTTAGTCACTGGC
Allele-1: 1bp insertion in exon2

WT CCATGGAACCAAGACAGAAAAAGCGGCTGTTAGTCACTGGCA
Mut CCATGGAACCAAGACAGAAAAAGCGGCTGTTAGTCACTGGCA
Allele-2: WT

Genome sequence analysis of PCR products from parental (WT) and CTNNB1 knockdown (KD) 293T cells, using sanger sequencing.