

SQSTM1 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50134

Basic Information

Catalog No.

RM50134

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

SQSTM1

Species

Human

Gene ID


8878

Swiss Prot

Q13501

Synonymsp60; p62; A170; DMRV; OSIL; PDB3;
ZIP3; p62B; NADGP; FTDALS3;
SQSTM1/p62

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF- κ B) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF- κ B in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone.

Product Information

Description

SQSTM1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:1bp insertion and 3bp deletion in exon2

Allele-2:1bp insertion and 17bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount50 μ L, 2 μ g/ μ L.**Storage**

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

ProtocolTo be used as WB control. Lysate is supplied in 1 \times SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT GCAA CAT**CCAGAC*****GACTTGTAGCGT
Mut GCAAACAT**CCAGAC***Deletion***GACTTGTAGCGT
Allele-1: 1bp Insertion and 3bp deletion in exon2

WT GCAA CAT**GTGCAG*****ACTTGTAGCGTC
Mut GCAAACAT**GTGCAG***Deletion***ACTTGTAGCGTC
Allele-2: 1bp Insertion and 17bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and SQSTM1 knockout (KO) 293T cells, using sanger sequencing.