

RIPK3 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM50138

Basic Information

Catalog No.

RM50138

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

RIPK3

Species

Human

Gene ID

11035

Swiss Prot

Q9Y572


Synonyms

RIP3; RIPK3

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

The product of this gene is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is predominantly localized to the cytoplasm, and can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor.

Product Information

Description

RIPK3 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:65bp deletion in exon2

Allele-2:65bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50µL, 2µg/µL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT CCCCTTGGTGCCA*****CATAGGAAGTGGGG
Mut CCCCTTGGTGCCA***Deletion***CATAGGAAGTGGGG
Allele-1: 65bp deletion in exon2
WT CCCCTTGGTGCCA*****CATAGGAAGTGGGG
Mut CCCCTTGGTGCCA***Deletion***CATAGGAAGTGGGG
Allele-2: 65bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and RIPK3 knockout (KO) HeLa cells, using sanger sequencing.