# ABclonal www.abclonal.com

# CHD1 Knockout 293T cell line, Homozygous

Catalog No.: RM50171

#### **Basic Information**

## Catalog No.

RM50171

#### Category

Cell Lysate

#### **Parental Cell line**

293T

#### Genotype

Knockout

## **Background**

#### **Gene Information**

#### **Gene Symbol**

CHD1

## **Species**

Human

## **Gene ID**

1105

#### **Swiss Prot**

014646

#### **Synonyms**

CHD-1; PILBOS; D1

#### **Contact**

2	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

## **Product Information**

#### **Description**

CHD1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:67bp deletion in exon2

Allele-2:67bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

## **Packaging**

1 vial parental cell line and 1 vial knockout cell line

## **Shipping Conditions**

**Amount** 

Dry ice

1~5x10<sup>6</sup> cells/vial.

#### **Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

#### **Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at  $37^{\circ}C$  with 5% CO<sub>2</sub> condition.

- 1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

WT ACTCCGGATCTGAA\*\*\*\*\*\*\*\*\*\*AAGTTGATGGAGCT Mut ACTCCGGATCTGAA\*\*\*Deletion\*\*\*AAGTTGATGGAGCT Allele-1: 67bp deletion in exon2

WT ACTCCGGATCTGAA\*\*\*\*\*\*\*\*\*AAGTTGATGGAGCT
Mut ACTCCGGATCTGAA\*\*Deletion\*\*\*AAGTTGATGGAGCT
Allele-2: 67bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and CHD1 knockout (KO) 293T cells, using sanger sequencing.