

TWIST1 Knockout 293T cell lysate, Homozygous

Catalog No.: RM50177

Basic Information

Catalog No.

RM50177

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

TWIST1

Species

Human

Gene ID

7291

Swiss Prot

Q15672

SynonymsCRS; CSO; SCS; ACS3; CRS1; BPES2;
BPES3; SWCOS; TWIST; bHLHa38; Twist

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a basic helix-loop-helix (bHLH) transcription factor that plays an important role in embryonic development. The encoded protein forms both homodimers and heterodimers that bind to DNA E box sequences and regulate the transcription of genes involved in cranial suture closure during skull development. This protein may also regulate neural tube closure, limb development and brown fat metabolism. This gene is hypermethylated and overexpressed in multiple human cancers, and the encoded protein promotes tumor cell invasion and metastasis, as well as metastatic recurrence. Mutations in this gene cause Saethre-Chotzen syndrome in human patients, which is characterized by craniosynostosis, ptosis and hypertelorism.

Product Information

Description

TWIST1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:122bp deletion in exon1

Allele-2:122bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT AGCGCGGGGACGC*****TGCGGGCTGTGGCG
Mut AGCGCGGGGACGC***Deletion***TGCGGGCTGTGGCG
Allele-1: 122bp deletion in exon1

WT AGCGCGGGGACGC*****TGCGGGCTGTGGCG
Mut AGCGCGGGGACGC***Deletion***TGCGGGCTGTGGCG
Allele-2: 122bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and TWIST1 knockout (KO) 293T cells, using sanger sequencing.