

Rabbit anti-Mouse CD278 mAb(DET)

Catalog No.: RMK0006

Basic Information

Catalog No.
RMK0006

Catagory
Elisa Antibody Kit

Application
ELISA,Multiplex

Product Information

Ig Type
Rabbit IgG

Purification
Affinity purification

Endotoxin Level

Storage
Store at -20°C. Avoid freeze / thaw cycles. Preservative 0.05%ProClin 300.
Avoid repeated freeze-thaw cycles.

Formulation
Supplied as a 0.2um filtered solution in PBS with 0.05%ProClin 300,PH 7.4.

Contact

 | order@abclonal.com

 | support@abclonal.com

 | www.abclonal.com

Background

Interleukin-10 (IL-10), also known as cytokine synthesis inhibitory factor (CSIF), is the charter member of the IL-10 alpha -helical cytokine family that also includes IL-19, IL-20, IL-22, IL-24, and IL-26/AK155 . IL-10 is secreted by many activated hematopoietic cell types as well as hepatic stellate cells, keratinocytes, and placental cytotrophoblasts. Whereas human IL-10 is active on mouse cells, mouse IL-10 does not act on human cells. Mature rat IL-10 shares 85% amino acid sequence identity with mouse IL-10 and 71% - 79% with bovine, canine, equine, feline, guinea pig, human, ovine, and porcine IL-10. It contains two intrachain disulfide bridges and is expressed as a 36 kDa noncovalently-associated homodimer.

Immunogen Information

Immunogen
Recombinant Mouse CD278

Cross-Reactivity


Assay Applications

Mouse CD278 Multiplex/ELISA


		Recommended Concentration	Sample
Multiplex	Capture	3-20ug/mL	Rabbit anti- Mouse CD278 (Cat. No.RMK0005)
	Detection	0.017-2ug/mL	Rabbit anti- Mouse CD278 (Cat. No.RMK0006)
	Standard	6.86-5000pg/mL	Recombinant Mouse CD278

ELISA	Capture	1-4ug/mL	Rabbit anti- Mouse CD278 (Cat. No.RMK0005)
	Detection	0.05-0.0125ug/mL	Rabbit anti- Mouse CD278 (Cat. No.RMK0006)
	Standard	31.25-2000pg/mL	Recombinant Mouse CD278

Validation Data



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.