Rabbit anti-Mouse IFN-gamma mAb



Catalog No.: RMK0050

Basic Information

Catalog No. RMK0050

Catagory Elisa Antibody Kit

Application Multiplex

Product Information

Ig Type Rabbit IgG

Purification Affinity purification

Endotoxin Level

Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Avoid repeated freeze-thaw cycles.

Formulation

Supplied as a 0.2µm filtered solution in PBS with 0.05% Proclin300, PH 7.4.

Contact

-	order@abclonal.com
6	support@abclonal.com
€	www.abclonal.com

Background

Interferon-gamma (IFN-gamma), also known as type II or immune interferon, exerts a wide range of immunoregulatory activities and is considered to be the prototype proinflammatory cytokine . Mature rat IFN-gamma exists as a noncovalently linked homodimer of 20 - 25 kDa variably glycosylated subunits. It shares 86% amino acid sequence identity with mouse IFN-gamma and 37%-45% with bovine, canine, cotton rat, equine, feline, human, porcine, and rhesus IFN-gamma. IFN-gamma dimers bind to IFN-gamma RI (alpha subunits) which then interact with IFN-gamma RII (beta subunits) to form the functional receptor complex of two alpha and two beta subunits. Inclusion of IFN-gamma RII increases the binding affinity for ligand and the efficiency of signal transduction. IFN- gamma is produced by a variety of immune cells under inflammatory conditions, notably by T cells and NK cells. It plays a key role in host defense by promoting the development and activation of Th1 cells, chemoattraction and activation of monocytes and macrophages, upregulation o...

Immunogen Information

Immunogen

Recombinant Mouse IFN-gamma Protein


Cross-Reactivity

1


Assay Applications

Mouse IFN-gamma Multiplex

		Recommended Concentration	Sample
	Capture	0.1-5ug/mL	Rabbit anti-Mouse IFN-gamma mAb(Cat. No. RMK0050)
Multiplex	Detection	0.5-2ug/mL	Rabbit anti-Mouse/Rat IFN-gamma mAb (Cat. No. RMK0001)
	Standard	6.86-5000pg/mL	Recombinant Mouse IFN-gamma Protein (Cat. No.RP01070)



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.