

# Rabbit anti-Mouse G-CSF mAb (CAP)

Catalog No.: RMK0111

## Basic Information

**Catalog No.**  
RMK0111

**Catagory**  
Elisa Antibody Kit

**Application**  
ELISA

## Product Information

**Ig Type**  
Rabbit IgG

**Purification**  
Affinity purification

**Endotoxin Level**

**Storage**  
Store at -20°C.  
**Avoid repeated freeze-thaw cycles.**

**Formulation**  
Supplied as a 0.2 µm filtered solution in PBS with 0.05% Proclin 300, PH 7.4.

## Contact

 | order@abclonal.com

 | support@abclonal.com

 | www.abclonal.com

## Background

Mouse granulocyte-colony stimulating factor (G-CSF) is a 24-25 kDa monomeric glycoprotein that regulates the proliferation, differentiation and activation of hematopoietic cells. Mouse G-CSF cDNA encodes a 208 amino acid (aa) precursor protein with a 30 aa signal sequence that is proteolytically cleaved to form a 178 aa O-glycosylated mature protein containing two intrachain disulfide bridges. In humans, two distinct cDNA clones, encoding a 204 aa form and a minor alternatively spliced 207 aa form of G-CSF precursors, have been isolated. Mouse and human G-CSF are 76% identical at the aa sequence level and the two proteins show species cross-reactivity. G-CSF is produced primarily by monocytes and macrophages upon activation by endotoxin, TNF-α or IL-1. Other cell types, including fibroblasts, endothelial cells, astrocytes and bone marrow stroma cells, can also secrete G-CSF after activation. In addition, various tumor cells express G-CSF constitutively.

## Immunogen Information

**Immunogen**  
Recombinant mouse G-CSF


## Cross-Reactivity

## Assay Applications

Mouse G-CSF Sandwich ELISA Immunoassay

	Recommended Concentration	Sample
ELISA Capture	1-4 µg/mL	Rabbit anti-Mouse G-CSF (CAP)(Cat. No. RMK0111)
ELISA Detection	0.008-0.033 µg/mL	Rabbit anti-Mouse G-CSF (DET)(Cat. No. RMK0112)
Standard	7.8-500 pg/mL	Recombinant Mouse G-CSF Protein(Cat. No. RP00573)

# Validation Data



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.