# Rabbit anti-Mouse G-CSF mAb (DET)



Catalog No.: RMK0112

# **Basic Information**

## Catalog No.

RMK0112

#### Catagory

Elisa Antibody Kit

### **Application**

**ELISA** 

### **Product Information**

# Ig Type

Rabbit IgG

# **Purification**

Affinity purification

## **Endotoxin Level**

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Preservative 0.05% ProClin 300. Avoid repeated freeze-thaw cycles.

#### **Formulation**

Supplied as a 0.2um filtered solution in PBS with 0.05%ProClin 300.PH 7.4.

## Contact

<u> </u>	order@abclonal.com
	support@abclonal.com
	www.abclonal.com

# **Background**

Mouse granulocyte-colony stimulating factor (G-CSF) is a 24-25 kDa monomeric glycoprotein that regulates the proliferation, differentiation and activation of hematopoietic cells. Mouse G-CSF cDNA encodes a 208 amino acid (aa) precursor protein with a 30 aa signal sequence that is proteolytically cleaved to form a 178 aa O-glycosylated mature protein containing two intrachain disulfide bridges. In humans, two distinct cDNA clones, encoding a 204 aa form and a minor alternatively spliced 207 aa form of G-CSF precursors, have been isolated. Mouse and human G-CSF are 76% identical at the aa sequence level and the two proteins show species cross-reactivity. G-CSF is produced primarily by monocytes and macrophages upon activation by endotoxin, TNF- $\alpha$  or IL-1. Other cell types, including fibroblasts, endothelial cells, astrocytes and bone marrow stroma cells, can also secrete G-CSF after activation. In addition, various tumor cells express G-CSF constitutively.

# **Immunogen Information**

### **Immunogen**


Recombinant Mouse G-CSF

# **Cross-Reactivity**

# **Assay Applications**

Mouse G-CSF Sandwich ELISA Immunoassay

	Recommended Concentration	Sample
ELISA Capture	1-4ug/mL	Rabbit anti-Mouse G-CSF (CAP)(Cat. No.RMK0111)
ELISA Detection	0.008-0.033ug/mL	Rabbit anti-Mouse G-CSF (DET)(Cat. No.RMK0112)
Standard	7.8-500pg/mL	Recombinant Mouse G-CSF Protein(Cat. No.RP00573)



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.