

Anti- Mouse GM-CSF mAb (DET)

Catalog No.: RMK0307

Basic Information

Catalog No.

RMK0307

Catagory

Elisa Antibody Kit

Application

ELISA

Product Information

Ig Type

Rabbit IgG

Purification

Affinity purification

Endotoxin Level

Storage


Store at -20°C. Avoid freeze / thaw cycles. Preservative 0.05%ProClin 300.


Avoid repeated freeze-thaw cycles.

Formulation

Supplied as a 0.2um filtered solution in PBS with 0.05%ProClin 300,PH 7.4.

Contact

 | order@abclonal.com

 | support@abclonal.com

 | www.abclonal.com

Background

GM-CSF was initially characterized as a factor that can support the in vitro colony formation of granulocyte-macrophage progenitors. It is also a growth factor for erythroid, megakaryocyte, and eosinophil progenitors. GM-CSF is produced by a number of different cell types (including T cells, B cells, macrophages, mast cells, endothelial cells, fibroblasts, and adipocytes) in response to cytokine or inflammatory stimuli. On mature hematopoietic cells, GM-CSF is a survival factor for and activates the effector functions of granulocytes, monocytes/macrophages, and eosinophils. GM-CSF promotes a Th1 biased immune response, angiogenesis, allergic inflammation, and the development of autoimmunity. It shows clinical effectiveness in ameliorating chemotherapy-induced neutropenia, and GM-CSF transfected tumor cells are utilized as cancer vaccines. The 22 kDa glycosylated GM-CSF, similar to IL-3 and IL-5, is a cytokine with a core of four bundled alpha-helices. Mature mouse GM-CSF shares 49%-54% amino acid sequence identity with canine, fe...

Immunogen Information

Immunogen

Recombinant Mouse GM-CSF

Cross-Reactivity


/

Assay Applications

Mouse GM-CSF Sandwich ELISA Immunoassay


	Recommended Concentration	Sample
ELISA Capture	1-4 ug/mL	Rabbit anti-Mouse GM-CSF (CAP)(Cat. No.RMK0306)
ELISA Detection	0.001-0.004ug/mL	Rabbit anti-Mouse GM-CSF (DET)(Cat. No.RMK0307)
Standard	7.8-500pg/mL	Recombinant Mouse GM-CSF Protein

Validation Data




This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.

Specificity



No significant cross-reactivity or interference was observed.

Stability



Placed at 37°C for 3 days, the stability of the standard curve all conform to CV <10%.