

Rabbit anti-Human IgM mAb

Catalog No.: RMK0353

Basic Information

Catalog No.
RMK0353

Catagory
Elisa Antibody Kit

Application
Multiplex

Product Information

Ig Type
Rabbit IgG

Purification
Affinity purification

Endotoxin Level
/

Storage
Store at -20°C.
Avoid repeated freeze-thaw cycles.

Formulation
Supplied as a 0.2 µm filtered solution in PBS with 0.05% Proclin 300, pH 7.4.

Contact

 | order@abclonal.com

 | support@abclonal.com

 | www.abclonal.com

Background

Immunoglobulins (Ig) are the antigen recognition molecules of B cells. An Ig molecule is made up of 2 identical heavy chains and 2 identical light chains (see MIM 147200) joined by disulfide bonds so that each heavy chain is linked to a light chain and the 2 heavy chains are linked together. Each Ig heavy chain has an N-terminal variable (V) region containing the antigen-binding site and a C-terminal constant (C) region, encoded by an individual C region gene, that determines the isotype of the antibody and provides effector or signaling functions. The heavy chain V region is encoded by 1 each of 3 types of genes: V genes (see MIM 147070), joining (J) genes (see MIM 147010), and diversity (D) genes (see MIM 146910). The C region genes are clustered downstream of the V region genes within the heavy chain locus on chromosome 14. The IGHM gene encodes the C region of the mu heavy chain, which defines the IgM isotype. Naive B cells express the transmembrane forms of IgM and IgD (see IGHD; MIM 1471770) on their surface. During an antibody ...

Immunogen Information

Immunogen
Human IgM Protein

Cross-Reactivity

/

Assay Applications

Human IgM Sanwich Immunoassay

	Recommended Concentration	Sample
Multiplex Capture	3-20 µg/mL	Rabbit anti-Human IgM mAb (Cat. No. RMK0353)
Multiplex Detection	0.017-2 µg/mL	Rabbit anti-Human IgM mAb (Cat. No. RMK0354)
Standard	6.86-5000 pg/mL	Recombinant Human IgM Protein

Validation Data

Standard Curve Chart

