Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Rabbit anti-Human IL-15 mAb

Catalog No.: RMK0671

Basic Information

Catalog No.
RMK0671

Catagory
Elisa Antibody Kit

Application
ELISA

Product Information

Ig Type
Rabbit IgG

Purification
Affinity purification

Endotoxin Level
N/A

Storage
Store at -20℃
Avoid repeated freeze-thaw cycles.

Formulation
Supplied as a 0.2 μm filtered solution in PBS with 0.05% Proclin 300, pH 7.3.

Contact

order@abclonal.com

support@abclonal.com

www.abclonal.com

Background

The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Alternatively spliced transcript variants of this gene have been reported.

Immunogen Information

Immunogen
Recombinant Human IL-15 protein

Cross-Reactivity


Human

Assay Applications


Human IL-15 Sandwich Immunoassay

		Recommended Concentration	Sample
ELISA	Capture	2-8 μg/mL	Rabbit anti-Human IL-15 mAb (Cat. No. RMK0671)
	Detection	0.0125-0.05 μg/mL	Rabbit anti-Human IL-15 mAb (Cat. No. RMK0672)
	Standard	3.91-250 pg/mL	Recombinant Human IL-15 Protein (Cat. No. RP01236)


Validation Data



No significant cross-reactivity or interference was observed.



Placed at 37°C for 7 days, the stability of the standard curve all conform to CV <10%.



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.