Recombinant Human IL-18 Protein

www.abclonal.com

ABclonal

Catalog No.: RP00006 Recombinant

Sequence Information

Species Gene ID **Swiss Prot** Human 3606 014116

Tags No tag

Synonyms IL18;IGIF;IL-18;IL-1g;IL1F4

Product Information

Purification Source E. coli ≥ 95 % as

determined by SDS-PAGE.

Calculated MW Observed MW

18 kDa 18.22 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Background

Interleukin-18 (IL-18, also known as interferon-gamma inducing factor) is a proinflammatory cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL-18 activation is induced by infection or tissue damage and contributes to disease pathology in chronic inflammation. In the presence of IL-12 or IL-15, IL-18 enhances anti-viral Th1 immune responses by inducing IFN-gamma production and the cytolytic activity of CD8+ T cells and NK cells. In the absence of IL-12 or IL-15, however, IL-18 promotes production of the Th2 cytokines IL-4 and IL-13 by CD4+ T cells and basophils. In the presence of IL-1 beta or IL-23, IL-18 induces the antigen-independent production of IL-17 by gamma $\delta\,T$ cells and CD4+ T cells.IL-18 also promotes myeloid dendritic cell maturation and triggers neutrophil respiratory burst.In cancer, IL-18 exhibits diverse activities including enhancing anti-tumor immunity, inhibiting or promoting angiogenesis, and promoting tumor cell metastasis.

Basic Information

Description

Recombinant Human IL-18 Protein is produced by E. coli expression system. The target protein is expressed with sequence (Tyr37-Asp193) of human IL-18 (Accession #NP 001553.1).

Bio-Activity

1.Measured by its binding ability in a functional ELISA. Immobilized recombinant Human IL18 at 5 μg/mL (100μL/well) can bind recombinant Human IL18BPa with a linear range of 0.156-1.372 μg/mL.|2.Measured by its ability to induce IFN-gamma secretion by KG\1 human acute myelogenous leukemia cells in the presence of TNFalpha(Catalog: RP00001). The ED₅₀ for this effect is 1.44-5.76 ng/mL, corresponding to a specific activity of $1.7 \times 10^5 \sim 6.9 \times 10^5$ units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Validation Data



Recombinant Human IL-18 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human IL18 at $5\mu g/mL$ (100 $\mu L/well$) can bind recombinant Human IL18BPa with a linear range of 0.156-1.372 $\mu g/mL$.



Recombinant Human IL-18 induce IFN-gamma secretion by KG \square 1 human acute myelogenous leukemia cells in the presence of TNF-alpha(Catalog: RP00001). The ED $_{50}$ for this effect is 1.44-5.76 ng/mL, corresponding to a specific activity of $1.7\times10^5\sim6.9\times10^5$ units/mg.