

Catalog No.: RP00019 **Recombinant**

Species	Gene ID	Swiss Prot
Human	90865	O95760

No tag

IL33;C9orf26;DVS27;IL1F11;NF-
HEV:NFEHEV

Source <i>E. coli</i>	Purification ≥ 95 % as determined by SDS- PAGE
---------------------------------	--

17.99 kDa 19 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Interleukin 33 (IL-33), also known as DVS27 or NF-HEV (Nuclear Factor from High Endothelial venules), is a proinflammatory protein and a chromatin-associated cytokine of the IL-1 family with high sequence and structural similarity to IL-1 and IL-18. The encoded protein is involved in the maturation of Th2 cells and the activation of mast cells, basophils, eosinophils and natural killer cells.

Recombinant Human IL-33 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Ser112-Thr270) of human IL-33 (Accession #NP_001300973.1).


Measured by its binding ability in a functional ELISA. Immobilized Recombinant human IL33 at 10 µg/mL (100 µL/well) can bind Recombinant human IL1RL1 with a linear range of 37.8-151.4 ng/mL.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human IL-33 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant human IL33 at 10 $\mu\text{g/mL}$ (100 μL /well) can bind Recombinant human IL1RL1 with a linear range of 37.8-151.4 ng/mL.