

Catalog No.: RP00039 **Recombinant**

Species	Gene ID	Swiss Prot
Human	1270	P26441

C-His

CNTF:HCNTF

Source	Purification
<i>E. coli</i>	≥ 95 % as determined by SDS-PAGE

Calculated MW	Observed MW
23.77 kDa	27 kDa

< 1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μ m filtered solution of 20mM Tris, 150mM NaCl, pH 8.0. Contact us for customized product form or formulation.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

The protein is a polypeptide hormone whose actions appear to be restricted to the nervous system where it promotes neurotransmitter synthesis and neurite outgrowth in certain neuronal populations. The protein is a potent survival factor for neurons and oligodendrocytes and may be relevant in reducing tissue destruction during inflammatory attacks. A mutation in this protein, which results in aberrant splicing, leads to ciliary neurotrophic factor deficiency, but this phenotype is not causally related to neurologic disease.

Recombinant Human CNTF Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Met1-Met200) of human CNTF (Accession #NP_000605.1) fused with a 6×His tag at the C-terminus.


Measured in a cell proliferation assay using TF-1 human erythroleukemic cell line. The ED₅₀ for this effect is 13.5-54 ng/mL.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CNTF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Human CNTF stimulates cell proliferation of the TF-1 human erythroleukemic cell line. The ED_{50} for this effect is 13.5-54 ng/mL.