

Recombinant Human TIGIT Protein

Catalog No.: RP00080 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	201633	Q495A1

Tags

C-His

Synonyms

VSIG9; VSTM3; WUCAM;TIGIT

Product Information

Source	Purification
HEK293 cells	> 97% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

This protein is a member of the PVR (poliovirus receptor) family of immunoglobulin proteins. The protein is expressed on several classes of T cells including follicular B helper T cells (TFH). The protein has been shown to bind PVR with high affinity; this binding is thought to assist interactions between TFH and dendritic cells to regulate T cell dependent B cell responses.

Basic Information

Description

Recombinant Human TIGIT Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Met22-Pro141) of human TIGIT (Accession #NP_776160.2) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human TIGIT at 1.5 μg/mL (100 μL/well) can bind Human CD155 with a linear range of 0.15-1.2 μg/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human TIGIT Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 18-22 kDa.



Immobilized Human TIGIT at 1.5 µg/mL (100 µL/well) can bind Human CD155 with a linear range of 0.15-1.2 µg/mL.