

Recombinant Human IGFBP-6 Protein

Catalog No.: RP00145 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3489	P24592

Tags

C-His

Synonyms

IGFBP6;IBP6

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
26.16 kDa	35-40 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

The superfamily of insulin-like growth factor (IGF) binding proteins include the six high-affinity IGF binding proteins (IGFBP) and at least four additional low-affinity binding proteins referred to as IGFBP related proteins (IGFBP-rP). All IGFBP superfamily members are cysteine-rich proteins with conserved cysteine residues, they can bind IGF-I and IGF-II with the equal affinity. Insulin-like growth factor (IGF) binding proteins (IGFBPs) have been shown to either inhibit or enhance the action of IGF, or act in an IGF-independent manner in the prostate. IGF-binding protein-4 (IGFBP-4) inhibits IGF-I action in vitro and is the most abundant IGFBP in the rodent arterial wall. IGFBP6 is directly downregulated by the beta-catenin/TCF complex in desmoid tumors, and imply a role for the IGF axis in the proliferation of desmoid tumors. There is mounting evidence that the structure of the IGFBP proteins plays a key role in the regulation of IGF bioavailability, by modulating its molecular size, capillary membrane permeability, target tissue specificity, cell membrane adherence and IGF affinity.

Basic Information

Description

Recombinant Human IGFBP-6 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Gly240) of human IGFBP6/IBP-6 (Accession #NP_002169.1) fused with a 6xHis tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized recombinant human IGFBP6 at 1 μg/mL (100 μL/well) can bind recombinant human IGF1 with a linear range of 30-250 ng/mL.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Validation Data



Recombinant Human IGFBP-6 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 36 kDa.



The purity of Human IGFBP6/IBP-6 Protein (Cat.RP00145) was greater than 95% as determined by SEC-HPLC.



Immobilized recombinant human IGFBP6 at 1 $\mu\text{g/mL}$ (100 $\mu\text{L/well}$) can bind recombinant human IGF1 with a linear range of 30-250 ng/mL.