# Recombinant Human uPA/PLAU Protein

ABclonal

www.abclonal.com

Catalog No.: RP00146LQ Recombinant

## **Sequence Information**

**Species** Gene ID **Swiss Prot** Human 5328 P00749

**Tags** 

C-His

**Synonyms** 

PLAU; ATF; BDPLT5; QPD; UPA; URK; u-PA:urokinase

### **Product Information**

**Purification** HEK293 cells > 97% by SDS-

PAGE.

Calculated MW Observed MW

49.36 kDa 55 kDa

**Endotoxin** 

< 0.1 EU/µg of the protein by LAL method.

#### **Formulation**

Supplied as a 0.22 µm filtered solution in 20mM HEPES, 150mM NaCl, 2mM CaCl, 10% Glycerol, pH 7.5Contact us for customized product form or formulation.

### Reconstitution

### **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\overline{\mathfrak{S}}$	www.abclonal.com.cn

# **Background**

Plasminogen activator, urokinase, also known as PLAU and uPA, is a serine protease which converts plasminogen to plasmin, a broad-spectrum protease active on extracellular matrix (ECM) components. It is involved in complement activation, cell migration, wound healing, and generation of localized extracellular proteolysis during tissue remodelling, pro-hormone conversion, carcinogenesis and neoplasia. uPA and its receptor (uPAR) have been implicated in a broad spectrum of pathophysiological processes, including fibrinolysis, proteolysis, inflammation, atherogenesis and plaque destabilization, all of which are involved in the pathogenesis of MI (myocardial infarction). The role of uPA is not only does it as a kind of enzyme, but also is breast cancer, stomach cancer, colon cancer, bladder cancer, ovarian cancer, brain, and endometrial cancer markers for a strong invasion and metastasis. Because of the causal involvment of uPA in cancer invasion and metastasis, the blockade of uPA interactions and activity with specific inhibitors is of interest for novel strategies in cancer therapy.

### **Basic Information**

#### **Description**

Recombinant Human uPA/PLAU Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Leu431) of human Urokinase/PLAU (Accession #NP\_002649.1) fused with a 8×His tag at the C-terminus.


#### **Bio-Activity**

Measured by its binding ability in a functional ELISA. Immobilized Human uPAR Protein at 1 µg/mL (100 µL/well) can bind PLAU with a linear range of 0.031-1.014 ng/mL.


#### Storage

Store at -70°C. This product is stable at  $\leq$  -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.

# **Validation Data**



Recombinant Human uPA/PLAU Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55 kDa.



Immobilized Recombinant Human uPAR Protein at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind PLAU with a linear range of 0.031-1.014 ng/mL.