

Recombinant Human uPA/PLAU Protein

Catalog No.: RP00146LQ **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	5328	P00749

Tags

C-His

Synonyms

PLAU;ATF;BDPLT5;QPD;UPA;URK;u-PA;urokinase

Product Information

Source	Purification
HEK293 cells	> 97% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Supplied as a 0.22 μm filtered solution in 20mM HEPES, 150mM NaCl, 2mM CaCl, 10% Glycerol, pH 7.5Contact us for customized product form or formulation.

Reconstitution

Background

Plasminogen activator, urokinase, also known as PLAU and uPA, is a serine protease which converts plasminogen to plasmin, a broad-spectrum protease active on extracellular matrix (ECM) components. It is involved in complement activation, cell migration, wound healing, and generation of localized extracellular proteolysis during tissue remodelling, pro-hormone conversion, carcinogenesis and neoplasia. uPA and its receptor (uPAR) have been implicated in a broad spectrum of pathophysiological processes, including fibrinolysis, proteolysis, inflammation, atherogenesis and plaque destabilization, all of which are involved in the pathogenesis of MI (myocardial infarction). The role of uPA is not only does it as a kind of enzyme, but also is breast cancer, stomach cancer, colon cancer, bladder cancer, ovarian cancer, brain, and endometrial cancer markers for a strong invasion and metastasis. Because of the causal involvement of uPA in cancer invasion and metastasis, the blockade of uPA interactions and activity with specific inhibitors is of interest for novel strategies in cancer therapy.

Basic Information

Description

Recombinant Human uPA/PLAU Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Leu431) of human Urokinase/PLAU (Accession #NP_002649.1) fused with a 8×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human uPAR Protein at 1 μg/mL (100 μL/well) can bind PLAU with a linear range of 0.031-1.014 ng/mL.


Storage

This product is stable at ≤ -70° C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Validation Data



Recombinant Human uPA/PLAU Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55 kDa.



Immobilized Recombinant Human uPAR Protein at 1 μ g/mL (100 μ L/well) can bind PLAU with a linear range of 0.031-1.014 ng/mL.