

Recombinant Human B7-DC/PD-L2/CD273 Protein

Catalog No.: RP00150 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	80380	Q9BQ51

Tags

C-His

Synonyms

B7DC;Btdc;CD273;PD-L2;PDCD1L2;PDL2;bA574F11.2;PDCD1LG2

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
23.40 kDa	35-50 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

Programmed Death Ligand 2 (PD-L2), also known as B7-DC and butyrophilin-like protein, is a member of the B7 family of proteins that provide signals for regulating T-cell activation and tolerance. PD-L2 is expressed on dendritic cells, subsets of activated CD4+ and CD8+ T cells, and memory B cells that differentiate into plasma cells. At inflammatory sites such as rheumatoid arthritis, allergen exposure, and virus infection, PD-L2 is up-regulated on synoviocytes, infiltrating macrophages, dendritic cells, and airway epithelial cells. PD-L2, along with B7-H1/PD-L1, binds to T cell PD-1 where it promotes IFN-gamma production and CD40 Ligand up-regulation while inhibiting IL-4 production. In addition, PD-L2 binds to RGM-B on macrophages and alveolar epithelial cells, supporting respiratory immune tolerance. In asthma, PD-L2 suppresses IL-5 and IL-13 production, promotes IL-12 production by dendritic cells, and supports allergen-induced airway hyper-responsiveness and mucus production.

Basic Information

Description

Recombinant Human B7-DC/PD-L2/CD273 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Leu20-Pro219) of human B7-DC/PD-L2/CD273 (Accession #NP_079515.2) fused with a 6xHis tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human PD-L2 at 5 μg/mL (100 μL/well) can bind Recombinant Human PD-1 with a linear range of 0.3-1.2 μg/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Active Recombinant Human B7-DC/PD-L2/CD273 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 35-50 kDa.



The purity of Human B7-DC/PD-L2/CD273 Protein (Cat.RP00150) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Human PD-L2 at 5µg/mL (100 µL/well) can bind Recombinant Human PD-1 with a linear range of 0.3-1.2 µg/mL.