

Recombinant Human MHC class I polypeptide-related sequence A/MIC-A Protein

Catalog No.: RP00172 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	100507436	Q29983

Tags

C-His

Synonyms

MICA; MIC-A; PERB11.1; MHC class I polypeptide-related sequence A; MIC-A; PERB11.1

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
35.88 kDa	50-70 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

MHC class I chain-related molecules A (MICA) is one of the genes in the HLA class I region, which belongs to MHC class I family. It is the member of the non-classical class I family that displays the greatest degree of polymorphism. The MICA protein product is expressed on the cell surface, although unlike canonical class I molecules it does not seem to associate with beta-2-microglobulin. It is a ligand for the NKG2-D type II integral membrane protein receptor. The protein functions as a stress-induced antigen that is broadly recognized by intestinal epithelial gamma delta T cells. Variations in this gene have been associated with susceptibility to psoriasis 1 and psoriatic arthritis, and the shedding of MICA-related antibodies and ligands is involved in the progression from monoclonal gammopathy of undetermined significance to multiple myeloma.

Basic Information

Description

Recombinant Human MHC class I polypeptide-related sequence A/MIC-A Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Gln308) of human MICA (Accession #NP_000238.1) fused with a 6xHis tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human MICA Protein at 2 μg/mL (100 μL/well) can bind NKG2D with a linear range of 0.61-14.3 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human MIC-A Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55-65 kDa..



Immobilized Recombinant Human MICA Protein at 2 μ g/mL (100 μ L/well) can bind NKG2D with a linear range of 0.61-14.3 ng/mL.