

Recombinant Human IL-17A/CTLA-8 Protein

Catalog No.: RP00212 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3605	Q16552

Tags

C-His

Synonyms

IL17A; CTLA-8; CTLA8; IL-17; IL-17A;
IL17;
interleukin-17A;CTLA-8;CTLA8;IL-17;IL-17
A;IL17

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Calculated MW	Observed MW
16.37 kDa	18-25 kDa

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

IL17, also known as IL17a and CTLA8, is a 15-20 kDa glycosylated cytokine which belongs to the IL-17 family. The IL-17 family of cytokines includes six members, IL-17/IL-17A, IL-17B, IL-17C, IL-17D, IL-17E/IL-25, and IL-17F, which are produced by multiple cell types. IL17A promotes protective mucosal and epidermal inflammation in response to microbial infection. It induces chemokine production, neutrophil influx, and the production of antibacterial peptides. IL17A additionally enhances the production of inflammatory mediators by rheumatoid synovial fibroblasts and contributes to TNF alpha induced shock. In contrast, it can protect against the progression of colitis by limiting chronic inflammation. IL17A encourages the formation of autoreactive germinal centers and exacerbates the onset and progression of experimental models of autoimmunity.

Basic Information

Description

Recombinant Human IL-17A/CTLA-8 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ile20-Ala155) of human IL-17A (Accession #NP_002181.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant human IL17RA at 2 μg/mL (100 μL/well) can bind Recombinant human IL-17A, the EC₅₀ of human IL-17A is 32.5-130 ng/mL. 2. Measured by its ability to induce IL-6 secretion by Hela cells. The ED₅₀ for this effect is 2.64-10.58 ng/mL, corresponding to a specific activity of 9.45×10⁴~3.79×10⁵ units/mg.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Human IL-17A Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 18-25 kDa.



Immobilized Recombinant human IL17RA , His Tag at 2 µg/mL (100 µL/well) can bind Recombinant human IL-17A, The EC_{50} of human IL-17A is 32.5-130 ng/mL.



Recombinant Human IL-17A induce IL-6 secretion by Hela cells. The ED_{50} for this effect is 2.64-10.58 ng/mL, corresponding to a specific activity of $9.45 \times 10^4 \sim 3.79 \times 10^5$ units/mg.