

# Recombinant Human ICAM-1/CD54 Protein

Catalog No.: RP00272

Recombinant

2 Publications

## Sequence Information

Species	Gene ID	Swiss Prot
Human	3383	P05362

### Tags

C-His

### Synonyms

BB2;CD54;P3.58;ICAM1

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Background

This protein is a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor. ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation.

## Basic Information

### Description

Recombinant Human ICAM-1/CD54 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln28-Glu480) of human ICAM-1/CD54 (Accession #NP\_000192.2) fused with a 6×His tag at the C-terminus.

### Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized APC Mouse Anti-Human CD54 at 1μg/mL (25 μL/well) can bind Human CD54 with a linear range of 0.46-7.8 ng/mL.

### Storage

Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.


## Contact

☎ | 400-999-6126


✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

# Validation Data



Recombinant Human ICAM-1/CD54 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 70-80 kDa.



Immobilized APC Mouse Anti-Human CD54 at 1µg/mL (25 µL/well) can bind Human CD54 with a linear range of 0.46-7.8 ng/mL.