

# Recombinant Human NKAT-2/KIR2DL3/CD158b2 Protein

Catalog No.: RP00277 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	3804	AAB36590.1

### Tags

C-hFc&His

### Synonyms

KIR2DL3;CD158B2;CD158b;GL183;KIR-023GB;KIR-K7b;KIR-K7c;KIR2DS5;KIRCL23;NKAT;NKAT2;NKAT2A;NKAT2B;p58

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

## Background

## Basic Information

### Description

Recombinant Human NKAT-2/KIR2DL3/CD158b2 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (His22-His245) of human KIR2DL3/CD158b2 (Accession #NP\_056952.2) fused with an Fc, 6×His tag at the C-terminus.

### Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human KIR2DL3 at 1 μg/mL (100 μL/well) can bind KIR2DL3 Rabbit pAb with a linear range of 2-54 ng/mL.

### Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Contact

☎ | 400-999-6126


✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

Validation Data



Recombinant Human NKAT-2/KIR2DL3/CD158b2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 70-80 kDa.



Immobilized recombinant Human KIR2DL3 at 1 µg/mL (100 µL/well) can bind KIR2DL3 Rabbit pAb with a linear range of 2-54 ng/mL.