

Recombinant Human NKAT-2/KIR2DL3/CD158b2 Protein

Catalog No.: RP00277 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 3804 AAB36590.1

Tags

C-hFc&His

Synonyms

KIR2DL3;CD158B2;CD158b;GL183;KIR-02 3GB;KIR-K7b;KIR-K7c;KIR2DS5;KIRCL23;NKAT;NKAT2;NKAT 2A;NKAT2B;p58

Product Information

Source Purification
HEK293 cells > 95% by SDSPAGE.

Calculated MW Observed MW

51.38 kDa 70-80 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

| a | 400-999-6126 |
|-----------|---------------------------|
| \bowtie | cn.market@abclonal.com.cn |
| | www.abclonal.com.cn |

Background

Basic Information

Description

Recombinant Human NKAT-2/KIR2DL3/CD158b2 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (His22-His245) of human KIR2DL3/CD158b2 (Accession $\#NP_056952.2$) fused with an Fc, $6\times His$ tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA.Immobilized Human KIR2DL3 at 1 μ g/mL (100 μ L/well) can bind KIR2DL3 Rabbit pAb with a linear range of 2-54 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human NKAT-2/KIR2DL3/CD158b2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 70-80 kDa.



Immobilized recombinant Human KIR2DL3 at 1 μ g/mL (100 μ L/well) can bind KIR2DL3 Rabbit pAb with a linear range of 2-54 ng/mL.