

www.abclonal.com

## Recombinant

## Background

**Tags**  
C-His

CD262;DR5;KILLER;KILLER/DR5;TRAIL-  
R2;TRAILR2;TRICK2;TRICK2A;TRICK2B;TR  
ICKB;ZTNFR9;TNFRSF10B

This protein is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein.

### Description

Recombinant Human TNFRSF10B/DR5/TRAIL-R2/CD262 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Ile 56 - Glu 182) of human DR5 (Accession #NP\_003833.3) fused with a 6xHis tag at the C-terminus.

<b>Source</b>	<b>Purification</b>
HEK293 cells	≥ 90 % as determined by SDS-PAGE.

Calculated MW	Observed MW
15.13 kDa	17 kDa

## Endotoxin

< 0.1 EU/μg of the protein by LAL method.

## Formulation

Lyophilized from a 0.22  $\mu\text{m}$  filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

## Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human TRAIL at 2 µg/mL (100 µL/well) can bind Recombinant Human DR5 with a linear range of 14-56 ng/mL. 2. Measured by its ability to inhibit TRAIL-mediated cytotoxicity using L-929 mouse fibroblast cells treated with TRAIL. The ED<sub>50</sub> for this effect is 0.38-1.5 pg/mL in the presence of 20 ng/mL Recombinant Human TRAIL/TNFSF10. 3. Measured by its binding ability in a functional ELISA. Immobilized Human TRAIL / TNFSF10 Protein at 2 µg/mL (100 µL/well) can bind Human CD262 Protein with a linear range of 0.5-14.1 ng/mL.

## Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Contact

 | 400-999-6126


 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)


## Validation Data




Recombinant Human TNFRSF10B/DR5/TRAIL-R2/CD262 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant Human TRAIL at 2 $\mu$ g/mL (100  $\mu$ L/well) can bind Recombinant Human DR5 with a linear range of 14-56 ng/mL.



Recombinant Human TNFRSF10B inhibit TRAIL-mediated cytotoxicity using L-929 mouse fibroblast cells treated with TRAIL. The ED<sub>50</sub> for this effect is 0.38-1.5 pg/mL in the presence of 20 ng/mL Recombinant Human TRAIL/TNFSF10.



Immobilized Human TRAIL / TNFSF10 Protein at 2 $\mu$ g/mL (100  $\mu$ L/well) can bind Human CD262 Protein with a linear range of 0.5-14.1 ng/mL.