

Recombinant Human Glypican-1/GPC1 Protein

Catalog No.: RP00297 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 2817 P35052

Tags

C-His

Synonyms GPC1;glypican

Product Information

Source Purification HEK293 cells > 97% by SDS-PAGE.

Endotoxin

Please contact us for more information.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

| a | 400-999-6126 |
|-----------|---------------------------|
| \bowtie | cn.market@abclonal.com.cn |
| • | www.abclonal.com.cn |

Background

Cell surface heparan sulfate proteoglycans are composed of a membrane-associated protein core substituted with a variable number of heparan sulfate chains. Members of the glypican-related integral membrane proteoglycan family (GRIPS) contain a core protein anchored to the cytoplasmic membrane via a glycosyl phosphatidylinositol linkage. These proteins may play a role in the control of cell division and growth regulation.

Basic Information

Description

Recombinant Human Glypican-1/GPC1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp24-Gln527) of human Glypican 1 (Accession #NP 002072.1) fused with a 6×His tag at the C-terminus.

Bio-Activity


Measured by its binding ability in a functional ELISA. Immobilized Human FGF2 at 1 μ g/mL (100 μ L/well) can bind Human Glypican 1 with a linear range of 2-80 ng/mL.

Storage


Store the lyophilized protein at -20 $^{\circ}$ C to -80 $^{\circ}$ C for long term. After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Glypican-1/GPC1 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 60 kDa.



Immobilized recombinant Human FGF2 at 1 $\mu g/mL$ (100 $\mu L/well)$ can bind Human Glypican 1 with a linear range of 2-80ng/mL.