

# Recombinant Human Transthyretin/TTR Protein

Catalog No.: RP00298 **Recombinant**

## Sequence Information

| Species | Gene ID | Swiss Prot |
|---------|---------|------------|
| Human   | 7276    | P02766     |

### Tags

C-His

### Synonyms

CTS; CTS1; HEL111; HsT2651; PALB;  
TBPA;TTR;CTS1;HEL111;HsT2651;PALB;T  
BPA

## Product Information

| Source       | Purification           |
|--------------|------------------------|
| HEK293 cells | > 90% by SDS-<br>PAGE. |

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

### Basic Information

#### Description

Recombinant Human Transthyretin/TTR Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gly 21 - Glu 147 ) of human Transthyretin (Accession #NP\_000362.1) fused with a 6xHis tag at the C-terminus.


#### Bio-Activity

#### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

## Validation Data

---



Recombinant Human Transthyretin/TTR Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 16 kDa.



The purity of Human Transthyretin Protein (Cat.RP00298) was greater than 90% as determined by SEC-HPLC.