

# Recombinant Human SIRP-gamma/CD172g Protein

Catalog No.: RP00310 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	55423	Q9P1W8

### Tags

C-His&Avi

### Synonyms

CD172g; FLJ42230; SIRP beta 2; SIRP gamma; SIRP-B2; SIRPG; Cripto-1; bA77C3.1; SIRP  $\beta$  2

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

### Endotoxin

< 1 EU/ $\mu$ g of the protein by LAL method

### Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Signal regulatory protein gamma (SIRP gamma, designated CD172g), also called SIRP beta 2, is a monomeric 45-47 kDa type I transmembrane protein belonging to the SIRP/SHPS (CD172) family of the Ig superfamily. SIRP gamma is probable immunoglobulin-like cell surface receptor. On binding with CD47, mediates cell-cell adhesion. Engagement on T-cells by CD47 on antigen-presenting cells results in enhanced antigen-specific T-cell proliferation and costimulates T-cell activation.

## Basic Information

### Description

Recombinant Human SIRP-gamma/CD172g Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu29-Pro360) of Human SIRP-gamma/CD172g (Accession #Q9P1W8) fused with a C-His&Avi tag at the C-terminus.

### Bio-Activity


### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Validation Data



Human SIRP Gamma on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%..



The purity of Human SIRP Gamma is greater than 95% as determined by SEC-HPLC.



Immobilized Human SIRP Gamma, His Tag at 5 µg/mL (100 µL/well) on the plate. Dose response curve for Human CD47, hFc Tag with the EC<sub>50</sub> of 2.46 µg/mL determined by ELISA.