

Biotinylated Recombinant Human Siglec-8 Protein

Catalog No.: RP00334B **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	27181	Q9NYZ4

Tags

C-His&Avi

Synonyms

CDw329; MGC59785; SAF2; SAF2SAF-2; Siglec-8; SIGLEC8L; SIGLEC8

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

Siglec-8, also known as SAF, is an approximately 75 kDa transmembrane glycoprotein in the Siglec family of sialic acid-binding immune regulatory molecules. Mature human Siglec-8 consists of a 347 amino acid (aa) extracellular domain (ECD) with three Ig-like domains. Putative adhesion molecule that mediates sialic-acid dependent binding to red blood cells. Preferentially binds to alpha-2,3-linked sialic acid. Also binds to alpha-2,6-linked sialic acid.

Basic Information

Description

Biotinylated Recombinant Human Siglec-8 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met17-Ala363) of Human Siglec-8 (Accession #Q9NYZ4) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Biotinylated Human Siglec-8 on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.