

Recombinant Human TACTILE/CD96 Protein

Catalog No.: RP00346 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	10225	P40200-2

Tags

C-mFc

Synonyms

CD96 molecule; CD96; DKFZp667E2122;
TACTILE

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Calculated MW	Observed MW
79.8 kDa	120-160 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method


Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The receptors CD96 and TIGIT are expressed on the surface of T and natural killer (NK) cells, and recent studies suggest both play important inhibitory roles in immune function. CD96 has been shown to modulate immune cell activity in mice, with Cd96^{-/-} mice displaying hypersensitive NK-cell responses to immune challenge and significant tumor resistance. The counterbalance between the putative inhibitory CD96 and TIGIT receptors and the activating receptor, CD226, offers unique strategies for immunology drug development.

Basic Information

Description

Recombinant Human TACTILE/CD96 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val22-Met503) of Human TACTILE/CD96 (Accession #P40200-2) fused with a C-mFc tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Human CD96 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



Immobilized Human CD96, mFc Tag at 5 µg/mL (100 µL/Well) on the plate. Dose response curve for Human CD155, hFc Tag with the EC_{50} of 0.21 µg/mL determined by ELISA.



Immobilized Human CD96, mFc Tag at 2 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-CD96 Antibody, hFc Tag with the EC_{50} of 31.8ng/mL determined by ELISA (QC Test).