

# Recombinant Human Nectin-2/PVRL2/CD112 Protein

Catalog No.: RP00382 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	5819	Q92692-2

### Tags

C-hFc

### Synonyms

CD112; MPH; Nectin2; PRR2; PVRL2; HVEB; PVRR2

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Calculated MW	Observed MW
62.3 kDa	70-80 kDa

### Endotoxin

&lt; 1 EU/μg of the protein by LAL method

### Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Nectin-2 is an adhesion molecule that has been reported to play a role in tumor growth, metastasis and tumor angiogenesis. Nectin-2 expression in ovarian cancer may support tumor cell adhesion, leading to growth and lymph node metastasis. Effect of VEGF on Nectin-2 expression as well as permeability was investigated in HUVEC.

## Basic Information

### Description

Recombinant Human Nectin-2/PVRL2/CD112 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln32-Leu360) of Human Nectin-2/PVRL2/CD112 (Accession #Q92692-2) fused with a C-hFc tag at the C-terminus.

### Bio-Activity


#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Validation Data



Human Nectin-2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human Nectin-2 is greater than 95% as determined by SEC-HPLC.



Immobilized Human PVRIg, mFc Tag at 1 µg/mL (100 µL/well) on the plate. Dose response curve for Human Nectin-2, hFc Tag with the  $EC_{50}$  of 24.1ng/mL determined by ELISA.