

# Recombinant Human PD-1/PDCD1 Protein

Catalog No.: RP00403 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	5133	Q15116-1

### Tags

C-His

### Synonyms

PDCD1; PD1; CD279; SLEB2; PD-1

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

### Endotoxin

< 1 EU/μg of the protein by LAL method

### Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Programmed cell death protein 1, also known as PD-1 and CD279, is a protein found on the surface of cells that has a role in regulating the immune system's response to the cells of the human body by down-regulating the immune system and promoting self tolerance by suppressing T cell inflammatory activity.

## Basic Information

### Description

Recombinant Human PD-1/PDCD1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu25-Gln167) of Human PD-1/PDCD1 (Accession #Q15116-1) fused with a C-His tag at the C-terminus.

### Bio-Activity


#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Validation Data




Human PD-1 on Tris-Bis PAGE under reduced. The purity is greater than 95%.




The purity of Human PD-1 is greater than 95% as determined by SEC-HPLC.



Immobilized Human PD-1, His Tag at 1 µg/mL (100 µL/Well). Dose response curve for Anti-PD-1 Antibody, hFc Tag with the EC<sub>50</sub> of 6.9ng/mL determined by ELISA (QC Test).



Human PD-L2, hFc Tag captured on CM5 Chip via Protein A can bind Human PD-1, His Tag with an affinity constant of 0.21 µM as determined in SPR assay (Biacore T200).



Human PD-L1, hFc Tag captured on CM5 Chip via Protein A can bind Human PD-1, His Tag with an affinity constant of 2.20 µM as determined in SPR assay (Biacore T200).