

# Recombinant Human 5'-Nucleotidase/NT5E/CD73 Protein

Catalog No.: RP00410 Recombinant

## **Sequence Information**

**Species Gene ID Swiss Prot** Human 4907 P21589-1

**Tags** C-hFc

**Synonyms** 

CD73; NT5E; NTE; 5' nucleotidase (CD73); 5'-NT; 5-NT; E5NT; EC 3.1.3.5; eN; eNT; NT; NT55'-nucleotidase; NT5; NTE; CALJA

## **Product Information**

Source Purification

HEK293 cells ≥ 95 % as determined by SDS-PAGE;≥ 95 % as determined by

HPLC.

Calculated MW Observed MW

84.5 kDa 85-100 kDa

#### **Endotoxin**

< 1 EU/ $\mu g$  of the protein by LAL method.

## **Formulation**

Lyophilized from 0.22  $\mu m$  filtered solution in 20 mM Tris, 120 mM NaCl (pH 7.5). Normally 8% trehalose is added as protectant before lyophilization.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

### Contact

8	400-999-6126
$\times$	cn.market@abclonal.com.cn

## **Background**

CD73, also known as ecto-5'-nucleotidase, is an enzyme that in humans is encoded by the NT5E gene.CD73 commonly serves to convert AMP to adenosine.The enzyme consists of a dimer of 2 identical 70-kD subunits bound by a glycosyl phosphatidyl inositol linkage to the external face of the plasma membrane. The enzyme is used as a marker of lymphocyte differentiation. A? deficiency of CD73 occurs in a variety of immunodeficiency diseases.

## **Basic Information**

## **Description**

Recombinant Human 5'-Nucleotidase/NT5E/CD73 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Trp27-Lys547) of Human 5'-Nucleotidase/NT5E/CD73 (Accession #P21589-1) fused with a C-hFc tag at the C-terminus.

### **Bio-Activity**


#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## **Validation Data**



Recombinant Human 5'-Nucleotidase/NT5E/CD73 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Human CD73 is greater than 95% as determined by SEC-HPLC.



Immobilized Human CD73, hFc Tag at 2  $\mu$ g/mL (100  $\mu$ L/well) on the plate. Dose response curve for Biotinylated Anti-CD73 Antibody, hFc Tag with the EC $_{50}$  of 22.6ng/mL determined by ELISA (QC Test).