

Recombinant Human KLRB1/CD161 Protein

Catalog No.: RP00426 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3820	Q12918-1

Tags

C-hFc

Synonyms

KLRB1; CLEC5B; NKR-P1A; NKR-P1A;
HNKR-P1a; CD161; Ly59; NKR; NKR-P1;
NKR-P1; NKR-P1ANKR

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

CD161 (NKR-P1) is a lectin-like receptor present on NK cells and rare T-cell subsets. We have observed CD161 expression in some cases of T-cell prolymphocytic leukemia (T-PLL) and found it to be useful in follow-up and detection of disease after treatment.

Basic Information

Description

Recombinant Human KLRB1/CD161 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln67-Ser225) of Human KLRB1/CD161 (Accession #Q12918-1) fused with a C-hFc tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Human CD161 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human CD161 is greater than 93% as determined by SEC-HPLC.



Immobilized Human CD161, hFc Tag at 2 µg/mL (100 µL/Well) on the plate. Dose response curve for Biotinylated Anti-CD161 Antibody, hFc Tag with the EC₅₀ of 40.7ng/mL determined by ELISA.