

Biotinylated Recombinant Human CD117 Protein

Catalog No.: RP00434B **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3815	P10721-2

Tags

C-His&Avi

Synonyms

SCFR; PBT; CD117; ckit; c-kit; EC 2.7.10; EC 2.7.10.1; KIT

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

The c-kit proto-oncogen (CD 117) has been shown to be present in several cell types including normal and neoplastic hemopoietic cells. Among normal BM cells, CD117 expression has been found in about half of the CD34 precursors including progenitors committed to the erythroid, granulo-monocytic, and megakaryocytic cell lineages.

Basic Information

Description

Biotinylated Recombinant Human CD117 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln26-Pro520) of Human CD117 (Accession #P10721-2) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Biotinylated Human CD117 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human CD117 is greater than 95% as determined by SEC-HPLC.



Immobilized Human SCF, No Tag at 5 µg/mL (100 µL/well) on the plate. Dose response curve for Biotinylated Human CD117, His Tag with the EC_{50} of 6.8 µg/mL determined by ELISA.