

Biotinylated Recombinant Human Fc-gamma RIII beta/FCGR3B/CD16b (S36R, N82D, I106V) Protein

Catalog No.: RP00473BLQ **Recombinant**

Sequence Information

| Species | Gene ID | Swiss Prot |
|---------|---------|------------|
| Human | 2215 | O75015 |

Tags

C-His&Avi

Synonyms

CD16; Fc gamma RIIB; FCG3; FCGR3B;
FcγRIIB; FCRIIB; IGFR3; CD16b (NA2);
CD16B; FCG3B; FCG3; FCGR3; FCR-10

Product Information

| Source | Purification |
|--------------|------------------------------------|
| HEK293 cells | > 95% by SDS-PAGE > 95% by HPLC |

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Supplied as 0.22 μm filtered solution in PBS (pH 7.4).

Reconstitution

Background

Human Fc gamma RIIB/CD16b Protein is a receptor for the Fc region of immunoglobulins gamma. Low affinity receptor. Binds complexed or aggregated IgG and also monomeric IgG. Contrary to III-A, is not capable to mediate antibody-dependent cytotoxicity and phagocytosis. May serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.

Basic Information

Description

Biotinylated Recombinant Human Fc gamma RIII beta/FCGR3B/CD16b (S36R, N82D, I106V) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly17-Ser200) of Human Fc gamma RIII beta/FCGR3B/CD16b (S36R,N82D,I106V) (Accession #O75015) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity

Storage

Store at -70°C. This product is stable at ≤ -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Validation Data



Biotinylated Human Fc gamma RIIB (NA1) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human Fc gamma RIIB (NA1) is greater than 95% as determined by SEC-HPLC.



Biotinylated Human Fc gamma RIIB (NA1), His Tag captured on CM5 Chip via anti-his antibody can bind Rituximab with an affinity constant of 8.00 μ M as determined in SPR assay (Biacore T200).