

Recombinant Human Fc gamma RIIIA/CD16a (F176V) Protein

Catalog No.: RP00478 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	2214	AAH17865

Tags

C-His&Avi

Synonyms

Fc gamma RIIIA; FCGR3; FCGR3A; FCGR111; FcgRIIIA; FcR-10; CD16A; FCG3; IGFR3; FcRIIIa; FcRIII; CD16; IMD20

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method


Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Human Fc gamma RIIIA/CD16a Protein is a receptor for the Fc region of IgG. Binds complexed or aggregated IgG and also monomeric IgG. Mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis.

Basic Information

Description


Recombinant Human Fc gamma RIIIA/CD16a (F176V) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly17-Gln208(F176V)) of Human Fc gamma RIIIA/CD16a (Accession #AAH17865) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data




Human Fc gamma RIIIA (V176) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Fc gamma RIIIA (V176) is greater than 95% as determined by SEC-HPLC.



Immobilized Human Fc gamma RIIIA (V176), His Tag at 1 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-Fc gamma RIIIA Antibody, hFc Tag with the EC₅₀ of 23.3ng/mL determined by ELISA (QC test).



Human Fc gamma RIIIA (V176), His Tag captured on CM5 Chip via Anti-His Antibody can bind Herceptin, hFc Tag with an affinity constant of 67.3nM as determined in SPR assay (Biacore T200).