

# Biotinylated Recombinant Human Fc gamma RI/CD64 Protein

Catalog No.: RP00483B **Recombinant**

## Sequence Information

| Species | Gene ID | Swiss Prot |
|---------|---------|------------|
| Human   | 2209    | P12314-1   |

### Tags

C-His&Avi

### Synonyms

CD64; CD64a; Fc gamma RI; FCG1; Fc-gamma RIA; FCGR1; FcgRI; FCRI; FcRIA; FLJ18345; FCGR1A; IGFR1; Fcr; Fcgr; FCE1A; FcERI

## Product Information

| Source       | Purification                    |
|--------------|---------------------------------|
| HEK293 cells | > 95% by SDS-PAGE > 95% by HPLC |

### Endotoxin

< 1 EU/μg of the protein by LAL method

### Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc gamma? RI (also known as CD64), Fc gamma? RII (CD32), and Fc gamma? RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type.

## Basic Information

### Description

Biotinylated Recombinant Human Fc gamma RI/CD64 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln16-Pro288) of Human Fc gamma RI/CD64 (Accession #P12314-1) fused with a C-His&Avi tag at the C-terminus.

### Bio-Activity


#### Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Validation Data



Biotinylated Human Fc gamma RI on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human Fc gamma RI is greater than 95% as determined by SEC-HPLC.



Biotinylated Human Fc gamma RI, His-Avi Tag captured on CM5 Chip via Anti-His Antibody can bind Trastuzumab with an affinity constant of 0.91 nM as determined in SPR assay (Biacore T200).