Recombinant Human CD47 Protein

Catalog No.: RP00529 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	961	008722-1

Tags

No Tag

Synonyms

CD47 glycoprotein; CD47 molecule; CD47; IAP; OA3; MER6

Product Information

Source	Purification
HEK293 cells	≥ 95 % as
	determined by SDS
	PAGE;≥ 95 % as
	determined by
	HPLC.

Calculated MW Observed MW 13.72 kDa 35-48 kDa

Endotoxin

 $< 1 EU/\mu q$ of the protein by LAL method.

Formulation

Lyophilized from 0.22 um filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn

Background

CD47 (Cluster of Differentiation 47) also known as integrin associated protein (IAP) is a transmembrane protein that in humans is encoded by the CD47 gene. CD47 belongs to the immunoglobulin superfamily and partners with membrane integrins and also binds the ligands thrombospondin-1 (TSP-1) and signal-regulatory protein alpha (SIRPa).CD-47 acts as a don't eat me signal to macrophages of the immune system which has made it a potential therapeutic target in some cancers.

Basic Information

Description

Recombinant Human CD47 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln19-Pro139) of Human CD47 (Accession #Q08722-1) fused with No Tag.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.




www.abclonal.com.cn



Recombinant Human CD47 Protein was determined by Tris-Bis PAGE under reducing conditions.

The purity of Human CD47 is greater than 95% as determined by SEC-HPLC.



Immobilized Human CD47, No Tag at 5 μ g/mL (100 μ L/well) on the plate. Dose response curve for Human SIRP alpha, hFc Tag with the EC_{s0} of 0.24 μ g/mL determined by ELISA.