

Recombinant Human CD47 Protein

Catalog No.: RP00529 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	961	Q08722-1

Tags
No Tag

Synonyms
CD47 glycoprotein; CD47 molecule;
CD47; IAP; OA3; MER6

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Endotoxin
< 1 EU/μg of the protein by LAL method

Formulation
Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution
Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126
 | cn.market@abclonal.com.cn
 | www.abclonal.com.cn

Background

CD47 (Cluster of Differentiation 47) also known as integrin associated protein (IAP) is a transmembrane protein that in humans is encoded by the CD47 gene. CD47 belongs to the immunoglobulin superfamily and partners with membrane integrins and also binds the ligands thrombospondin-1 (TSP-1) and signal-regulatory protein alpha (SIRPα). CD-47 acts as a don't eat me signal to macrophages of the immune system which has made it a potential therapeutic target in some cancers.


Basic Information

Description
Recombinant Human CD47 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln19-Pro139) of Human CD47 (Accession #Q08722-1) fused with No Tag.


Bio-Activity

Storage
Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Human CD47 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human CD47 is greater than 95% as determined by SEC-HPLC.



Immobilized Human CD47, No Tag at 5 µg/mL (100 µL/well) on the plate. Dose response curve for Human SIRP alpha, hFc Tag with the EC_{50} of 0.24 µg/mL determined by ELISA.