## **Recombinant Human CD47 Protein**

Catalog No.: RP00531 Recombinant

## **Sequence Information**

Species	Gene ID	Swiss Prot
Human	961	Q08722-1

Tags

C-hFc

#### Synonyms

CD47 glycoprotein; CD47 molecule; CD47; IAP; OA3; MER6

## **Product Information**

Source	Purification
HEK293 cells	≥ 95 % as
	determined by SDS
	PAGE;≥ 95 % as
	determined by
	HPLC.

# Calculated MW Observed MW 40.5 kDa 55-65 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

#### Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn

## Background

CD47 (Cluster of Differentiation 47) also known as integrin associated protein (IAP) is a transmembrane protein that in humans is encoded by the CD47 gene. CD47 belongs to the immunoglobulin superfamily and partners with membrane integrins and also binds the ligands thrombospondin-1 (TSP-1) and signal-regulatory protein alpha (SIRP $\alpha$ ).CD-47 acts as a don't eat me signal to macrophages of the immune system which has made it a potential therapeutic target in some cancers.

## **Basic Information**

#### Description

Recombinant Human CD47 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln19-Pro139) of Human CD47 (Accession #Q08722-1) fused with a C-hFc tag at the C-terminus.

### **Bio-Activity**


### Storage

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles


Avoid repeated freeze/thaw cycles.



www.abclonal.com.cn




The purity of Human CD47 is greater than 95% as determined by SEC-HPLC.



Immobilized Human SIRP alpha, His Tag at 1  $\mu$ g/mL (100  $\mu$ L/Well) on the plate. Dose response curve for Human CD47, hFc Tag with the EC<sub>s0</sub> of 5.0ng/mL determined by ELISA (QC Test).





Human CD47, hFc Tag captured on Protein A chip via Anti-His Antibody can bind Human Sirpa V2, His Tag with an affinity constant of 13.8nM as determined in SPR assay (Biacore T200).