

ABclonal

www.abclonal.com

Recombinant Human BST-1/CD157 Protein

Catalog No.: RP00541 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 683 010588

Tags C-His

Synonyms BST1;CD157

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

29.95 kDa 35-45 kDa

Endotoxin

< 0.01 EU/ μ g of the protein by LAL method

Formulation

Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

The cluster of differentiation (CD) system is a glycosyl phosphatidylinositol anchored membrane protein thatbelongs to the CD38 family. It is generally used in immunophynotyping. CD157 was discovered in a bonemarrow stromal cell line where it facilitates pre-B-cell growth. CD157 is a bifunctional ectoenzyme that exhibits both ADP-ribosyl cyclase and cyclic ADP ribose hydrolase activities followed with CD38. It plays a role inrheumatoid arthritis (RA) due to its enhanced expression in RA-derived bone marrow stromal cell lines. Studies have shown that this protein have a role in predicted to function as a cell surface receptor and animmunoregulatory molecule.

Basic Information

Description

Recombinant Human BST-1/CD157 Protein is produced by Human Cells expression system. The target protein is expressed with sequence (Gly29-Lys292) of human CD157/BST1/ADP-ribosyl cyclase 2/Cyclic ADP-ribose hydrolase 2 (Accession #Q10588) fused with a 6×His tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human BST-1/CD157 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.