

Biotinylated Recombinant Human CD38 Protein

Catalog No.: RP00553B **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	952	P28907-1

Tags

C-His&Avi

Synonyms

ADPRC 1; T10; CD38; ADP-ribosyl cyclase 1; cADPr hydrolase 1; CD38 molecule; EC 2.4.99.20; EC 3.2.2.6

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

CD38 (cluster of differentiation 38), also known as cyclic ADP ribose hydrolase, is a transmembrane glycoprotein found on the surface of some immune cells including plasma cells, activated or immature T and B cells, monocytes, and natural killer cells. CD38 participates in cell adhesion, signal transduction and calcium signaling.

Basic Information

Description

Biotinylated Recombinant Human CD38 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val43-Ile300) of Human CD38 (Accession #P28907-1) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Biotinylated Human CD38 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human CD38 is greater than 95% as determined by SEC-HPLC.



Immobilized Biotinylated Human CD38, His Tag at 1 µg/mL (100 µL/well) on the streptavidin precoated plate(5 µg/mL). Dose response curve for Anti-CD38 Antibody, hFc Tag with the EC_{50} of 8.1ng/mL determined by ELISA.