

Biotinylated Recombinant Human B7-1/CD80 Protein

Catalog No.: RP00562B

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	941	P33681-1

Tags

C-His&Avi

Synonyms

CD80 molecule; B7; B71; B7-1; BB1; CD28LG1; CD80; B7.1; CD28LG1; LAB7

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC.

Calculated MW	Observed MW
26.8 kDa	50-70 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎	400-999-6126
✉	cn.market@abclonal.com.cn

Background

Cluster of differentiation 80 (also CD80 and B7-1) is a protein found on dendritic cells, activated B cells and monocytes that provides a costimulatory signal necessary for T cell activation and survival. It is the ligand for two different proteins on the T cell surface: CD28 and CTLA-4.

Basic Information

Description

Biotinylated Recombinant Human B7-1/CD80 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val35-Asn242) of Human B7-1/CD80 (Accession #P33681-1) fused with a C-His&Avi tag at the C-terminus.


Bio-Activity

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.




Validation Data



Biotinylated Recombinant Human B7-1/CD80 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Biotinylated Human B7-1 is greater than 95% as determined by SEC-HPLC.



Immobilized Biotinylated Human B7-1, His Tag at 0.5 µg/mL (100 µL/Well) on streptavidin (5 µg/mL) precoated plate. Dose response curve for Human CTLA-4, hFc Tag with the EC₅₀ of 1.8ng/mL determined by ELISA.