

Biotinylated Recombinant Human MS4A1/CD20 Protein www.abclonal.com

Catalog No.: RP00584BLQ Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 931 P11836-1

Tags

C-His&Avi

Synonyms

MS4A1; CD20; MS4A-1; B1; Bp35; CD20 receptor; CD20S7; CVID5; LEU-16; Ly-44;

MS4A2; S7

Product Information

Source Purification *E. coli* > 95% by SDS-PAGE

Calculated MW Observed MW

16.8 kDa 20-25 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method

Formulation

Supplied as 0.22 µm filtered solution in 20 mM PB, 10% Glycerol (pH 7.0).

Reconstitution

Background

B-lymphocyte antigen CD20 or CD20 is an activated-glycosylated phosphoprotein expressed on the surface of all B-cells beginning at the pro-B phase (CD45R , CD117) and progressively increasing in concentration until maturity.CD20 is the target of the monoclonal antibodies rituximab, ocrelizumab, obinutuzumab, ofatumumab, ibritumomab tiuxetan, tositumomab, and ublituximab, which are all active agents in the treatment of all B cell lymphomas, leukemias, and B cell-mediated autoimmune diseases.

Basic Information

Description

Biotinylated Recombinant Human MS4A1/CD20 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Ile141-Ser188) of Human MS4A1/CD20 (Accession #P11836-1) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity


Storage

Store at -70°C. This product is stable at \leq -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.


Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
<u></u>	ī	www.abclonal.com.cn

Validation Data



Biotinylated Human CD20 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



Immobilized Biotinylated Human CD20, His Tag at 1 μ g/mL (100 μ L/well) on the streptavidin precoated plate (5 μ g/mL). Dose response curve for Anti-CD20 Antibody, hFc Tag with the EC₅₀ of 0.12 μ g/mL determined by ELISA.