# **Recombinant Human CD24 Protein-VLP**



Catalog No.: RP00590 Recombinant

# **Sequence Information**

**Species Gene ID Swiss Prot** Human 100133941 P25063-1

**Tags** No Tag

**Synonyms** 

CD 24; CD24 molecule; CD24A; FLJ22950; FLJ43543; MGC75043

# **Product Information**

**Source** Purification HEK293 cells

#### **Endotoxin**

< 1 EU/ $\mu g$  of the protein by LAL method

## **Formulation**

Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

#### **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Background**

CD24 is a sialoglycoprotein expressed at the surface of most B lymphocytes and differentiating neuroblasts. It is also expressed on neutrophils and neutrophil precursors from the myelocyte stage onwards. The potential for targeting CD24 in cancer therapy seems promising, as CD24 is overexpressed in many human cancers.

## **Basic Information**

#### **Description**

Recombinant Human CD24 Protein-VLP is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser27-Gly59) of Human CD24-VLP (Accession #P25063-1) fused with No Tag.


## **Bio-Activity**

#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.



Immobilized Human CD24 VLP at 0.5  $\mu$ g/mL (100  $\mu$ L/Well) on the plate. Dose response curve for Anti-CD24 Antibody, hFc Tag with the EC $_{50}$  of 16.4ng/mL determined by ELISA.