

# Biotinylated Recombinant Human CD3E&CD3G Protein

**Catalog No.: RP00591B** Recombinant

## Sequence Information

Species	Gene ID	Swiss Prot
Human	916&917	P07766(CD3E)&P09693(CD3G)

### Tags

C-hFc&amp;Avi

### Synonyms

 CD3; CD3G; CD3g; T3G; CD3e; CD3E;  
 CD3 epsilon&CD3 gamma; CD3E&CD3G

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

### Calculated MW Observed MW

 40 kDa (CD3E), 36.5 kDa (CD3G)  
 48-55 kDa (CD3G)

### Endotoxin

&lt; 1 EU/μg of the protein by LAL method

### Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Background

T-cell surface glycoprotein CD3 epsilon&CD3 gamma chain, also known as CD3E&CD3G, are single-pass type I membrane proteins. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain.

## Basic Information

### Description

Biotinylated Recombinant Human CD3E&CD3G/CD3 epsilon&CD3 gamma Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp23-Asp126(CD3E) & Gln23-Ser116(CD3G)) of Human CD3E&CD3G/CD3 epsilon&CD3 gamma (Accession #P07766(CD3E)&P09693(CD3G)) fused with a C-hFc&Avi tag at the C-terminus.

### Bio-Activity

### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.




| [www.abclonal.com.cn](http://www.abclonal.com.cn)

---


## Validation Data



Biotinylated CD3E&CD3G on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human CD3E&CD3G is greater than 95% as determined by SEC-HPLC.



Immobilized Biotinylated Human CD3E&CD3G, hFc Tag at 1 µg/mL (100 µL/Well) on streptavidin (5 µg/mL) precoated plate. Dose response curve for OKT3, mFc Tag with the EC<sub>50</sub> of 30.3ng/mL determined by ELISA.