

# Recombinant Human CD5 Protein

Catalog No.: RP00594 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	921	P06127

### Tags

C-hFc&Avi

### Synonyms

CD5 molecule; CD5; LEU1; T1; CD5 antigen

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE > 95% by HPLC

**Calculated MW**   **Observed MW**

### Endotoxin

< 1 EU/μg of the protein by LAL method

### Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

CD5: a type I transmembrane protein found on T cells, thymocytes, and some B cells that is a ligand for CD72 and is involved in cellular activation or adhesion; expressed in B-cell chronic lymphocytic leukemia and T-cell lymphoma.

## Basic Information

### Description

Recombinant Human CD5 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Arg25-Asn371) of Human CD5 (Accession #P06127) fused with a C-hFc&Avi tag at the C-terminus.

### Bio-Activity


### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Validation Data



Human CD5 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human CD5 is greater than 95% as determined by SEC-HPLC.



Immobilized Human CD5 at 0.5 µg/mL (100 µL/Well) on the plate. Dose response curve for Biotinylated Anti-CD5 Antibody, hFc Tag with the EC<sub>50</sub> 14.6ng/mL determined by ELISA.