ABclonal www.abclonal.com

Recombinant Human CD4/LEU3 Protein

Catalog No.: RP00596 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 920 P01730

Tags

C-His&Avi

Synonyms

CD4 antigen (p55); CD4 molecule; CD4 receptor; CD4; CD4mut; EU3

Product Information

Source Purification HEK293 cells > 95% by SDS-

PAGE[]> 95% by

Endotoxin

< 1 EU/µg of the protein by LAL method

Formulation

Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

CD4, also known as L3T4, T4, and W3/25, is an approximately 55 kDa type I transmembrane glycoprotein that is expressed predominantly on thymocytes and a subset of mature T lymphocytes. It is a standard phenotype marker for the identification of T cell populations.Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex.

Basic Information

Description

Recombinant Human CD4/LEU3 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Lys26-Trp390) of Human CD4/LEU3 (Accession #P01730) fused with a C-His&Avi tag at the C-terminus.


Bio-Activity

Storage


Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Human CD4 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human CD4 is greater than 95% as determined by SEC-HPLC



Immobilized Human CD4 at 0.5 μ g/mL (100 μ L/Well) on the plate. Dose response curve for Anti-CD4 Antibody, hFc Tag with the EC₅₀ 10.6ng/mL determined by ELISA.